

THE GROMOV WIDTH OF 4-DIMENSIONAL TORI

JANKO LATSCHEV, DUSA MCDUFF, AND FELIX SCHLENK

ABSTRACT. Let ω be any linear symplectic form on the 4-torus T^4 . We show that in all cases (T^4, ω) can be fully filled by one symplectic ball. If (T^4, ω) is not symplectomorphic to a product $T^2(\mu) \times T^2(\mu)$ of equal sized factors, then it can also be fully filled by one symplectic ellipsoid of any shape, or, more generally, by any finite collection of balls or ellipsoids provided only that their total volume is less than that of (T^4, ω) .

CONTENTS

1. Introduction	1
2. Relations to algebraic geometry	6
2.1. Basics	6
2.2. Seshadri constants of tori	7
2.3. Seshadri estimates for higher dimensional tori	10
3. Proof of Theorems 1.9 and 1.10	11
3.1. Irrational case	11
3.2. Irrational tori with no curves	12
3.3. Product tori	16
4. Basic symplectic mappings	17
4.1. Diamonds	17
4.2. Distorted diamonds	18
4.3. Shears	20
5. Proof of Theorem 1	25
6. Proofs of Corollaries 1.2 and 1.3	26
7. Remarks and Questions	27
References	29

1. INTRODUCTION

Time and again, tori have proven to be interesting and important test cases for questions in symplectic topology. In this work we compute the Gromov width of 4-dimensional tori with

Date: November 29, 2011.

2000 Mathematics Subject Classification. 53D05, 14B05, 32S05, 11A55.

Key words and phrases. Gromov width, symplectic embeddings, symplectic packing, symplectic filling, tori.

partially supported by NSF grant DMS 0905191.

partially supported by SNF grant 200021-125352/1.

linear symplectic forms. Consider the open ball of capacity a ,

$$B^{2n}(a) = \left\{ z \in \mathbb{C}^n \mid \pi \sum_{j=1}^n |z_j|^2 < a \right\},$$

in standard symplectic space $(\mathbb{R}^{2n}, \omega_0)$, where $\omega_0 = \sum_{j=1}^n dx_j \wedge dy_j$. The *Gromov width* of a $2n$ -dimensional symplectic manifold (M, ω) , introduced in [15], is defined as

$$(1.1) \quad c_G(M, \omega) = \sup \{ a \mid B^{2n}(a) \text{ symplectically embeds into } (M, \omega) \}.$$

Computations and estimates of the Gromov width for various examples can be found in [4, 5, 6, 15, 18, 19, 21, 22, 29, 34, 38, 41, 42].

If the symplectic manifold (M, ω) has finite volume, an invariant equivalent to its Gromov width is the *ball filling number*

$$p(M, \omega) = \sup \frac{\text{Vol}(B^{2n}(a))}{\text{Vol}(M, \omega)}$$

where the supremum is taken over all balls $B^{2n}(a)$ that symplectically embed into (M, ω) , and where the volume is defined as $\frac{1}{n!} \int_M \omega^n$. Since $\text{Vol}(B^{2n}(a)) = \frac{a^n}{n!}$,

$$(1.2) \quad p(M, \omega) = \frac{(c_G(M, \omega))^n}{n! \text{Vol}(M, \omega)}.$$

If $p(M, \omega) < 1$ one says that *there is a filling obstruction*, while if $p(M, \omega) = 1$ one says that (M, ω) *admits a full filling by one ball*.¹ In this paper our main focus is the filling number of 4-tori with a linear symplectic form ω , i.e. those which can be identified with the quotient of $\mathbb{R}^4$, with its standard symplectic structure, by a suitable lattice Λ . We also study other related filling questions in which the ball is replaced by a disjoint union of balls or ellipsoids.

Filling obstructions usually come from non-constant holomorphic spheres. In tori, however, there are no such spheres. One can thus believe that for tori there should be no filling obstructions. For the standard torus $T(1, 1) := \mathbb{R}^4/\mathbb{Z}^4$, there is the obvious lower bound $p(T(1, 1)) \geq \frac{1}{2}$ coming from the inclusion of the ball $B^4(1)$ into the polydisc $B^2(1) \times B^2(1)$; see also Figure 3 below. A better lower bound (namely $p(T(1, 1)) \geq \frac{8}{9}$) comes from algebraic geometry. As we explain in §2.2 (cf. equation (2.5) in particular), we can fill up to 8/9ths of the volume of $(T^4, \omega) = T(1, 1)$ by a ball that is embedded both symplectically and holomorphically² with respect to some complex structure with Kähler form ω .

Our main result is

Theorem 1.1. *Every 4-dimensional linear symplectic torus admits a full filling by one ball; in other words, $p(T^4, \omega) = 1$ for all linear ω .*

The symplectic (resp. Kähler) cone of a smooth oriented manifold X is the set of cohomology classes $\alpha \in H^2(X; \mathbb{R})$ that can be represented by a symplectic (resp. Kähler) form, where here we consider symplectic forms that are compatible with the given orientation on X (resp. Kähler forms that are compatible with any complex structure giving this orientation). The symplectic cone $\mathcal{C}(T^4)$ of T^4 with a given orientation is $\{\alpha \in H^2(T^4; \mathbb{R}) \mid \alpha^2 > 0\}$. Each such

¹Our *ball filling number* is called *first packing number* by other authors, and *full fillings by one ball* also go under the name of *full packings by one ball*. We refer to §7 for a discussion of *full fillings* versus *very full fillings*.

²Such a ball is embedded isometrically with respect to the corresponding Kähler metric.

class has a linear representative. From Theorem 1.1, we get the following characterization of the symplectic cone of the 1-point blow up of a given oriented torus T^4 .

Corollary 1.2. *Denoting by $E \in H_2(X_1; \mathbb{Z})$ the homology class of the exceptional divisor (with some orientation) in $X_1 = T^4 \# \overline{\mathbb{C}P^2}$, the symplectic cone of X_1 is*

$$\mathcal{C}(X_1) = \{ \alpha \in H^2(X_1; \mathbb{R}) \mid \alpha^2 > 0, \alpha(E) \neq 0 \}.$$

While there are many examples of non-Kähler symplectic manifolds, it is much harder to find Kähler manifolds for which the Kähler and symplectic cones differ. Some examples are given by Drăghici [13] and Li–Usher [28]. With the help of Corollary 1.2 we obtain simpler examples.

Corollary 1.3. *Let X_k be the k -fold blow-up $T^4 \#_k \overline{\mathbb{C}P^2}$ of the 4-torus by $k \geq 1$ points. Then the symplectic cone of X_k is strictly bigger than the Kähler cone.*

This holds because Kähler forms on the blow-up correspond to balls in the torus that are embedded both symplectically and holomorphically, and known results in algebraic geometry (cf. §2.2) give restrictions on the size of such balls.

Outline of the proof of Theorem 1.1

As we will see, for our purposes linear 4-tori divide into three classes: the *standard torus* $T(1, 1)$ (and its rescalings), *all other rational tori* (in which $[\omega]$ is a multiple of a rational class), and *irrational tori* (in which the image of the homomorphism $\int \omega: H_2(M; \mathbb{Z}) \rightarrow \mathbb{R}$ has rank at least 2).

It turns out that every rational torus is (up to scaling) symplectomorphic to a product torus $T^2(d_1) \times T^2(d_2)$, where $d_1, d_2 \in \mathbb{N}$ denote the areas of the two factors; see Lemma 2.1. Thus the family $T(1, \mu) := T^2(1) \times T^2(\mu)$ of product tori contains all rational tori, up to scaling. With this in mind, our proof proceeds as follows.

1. *Linear algebra.* By a simple symplectic linear algebra argument, the tori $T(m, n)$ and $T(1, mn)$ are symplectomorphic for relatively prime integers m, n (see Remark 2.2). Hence:

Lemma 1.4. $p(T(\frac{m}{n}, 1)) = p(T(1, mn))$ for $m, n \in \mathbb{N}$ relatively prime.

2. *Algebraic geometry.* Buchdahl [8] and Lamari [24] found a condition on a cohomology class $\alpha \in H^{1,1}(X; \mathbb{R})$ on some complex surface X with even first Betti number that guarantees the existence of a Kähler representative of α . We shall verify this condition on blow-ups of irrational tori to obtain:

Proposition 1.5. $p(T^4, \omega) = 1$ for all irrational linear tori (T^4, ω) .

3. *Full fillings of $T^2(1) \times S^2(\mu)$.* Denote by $S^2(\mu)$ the 2-sphere endowed with an area form of area μ . Biran [4] proved that $T^2(1) \times S^2(\mu)$ can be fully packed by one ball provided that $\mu \geq 2$. We shall show that such an almost filling ball can be made to lie in the complement of a constant section $T^2(1) \times \text{pt}$. Since the open disc bundle $T^2(1) \times D^2(\mu) = (T^2(1) \times S^2(\mu)) \setminus (T^2(1) \times \text{pt})$ symplectically embeds into $T^2(1) \times T^2(\mu) = T(1, \mu)$, we obtain

Proposition 1.6. $p(T(\mu, 1)) = 1$ for all $\mu \geq 2$.

Corollary 1.7. $p(T(\mu, 1)) = 1$ for all $\mu \neq 1$.

Proof. In view of Propositions 1.5 and 1.6 we need only consider $\mu \in (1, 2) \cap \mathbb{Q}$. If we write $\mu = \frac{m}{n}$ with $m, n \in \mathbb{N}$ relatively prime, then $mn \geq 6$. Hence Lemma 1.4 and Proposition 1.6 imply $p(T(\mu, 1)) = p(T(1, mn)) = 1$. $\square$

4. *A symplectic embedding construction.* The only case not covered by the above discussion is the standard product torus $T(1, 1)$. To prove $p(T(1, 1)) = 1$ we shall construct for each ball $B^4(a)$ of volume $\frac{a^2}{2} < 1$ an explicit symplectic embedding into $T(1, 1)$. Fix $a < \sqrt{2}$. We start with an almost full embedding $B^4(a) \rightarrow \diamond \times \square$, where $\diamond \subset \mathbb{R}^2(x_1, x_2)$ is a diamond-shaped domain (see Figure 3 (I) below), and $\square = (0, 1)^2 \subset \mathbb{R}^2(y_1, y_2)$. The main step is then to construct a symplectic embedding $\diamond \times \square \rightarrow \mathbb{R}^4$ with image U such that the projection $\mathbb{R}^4 \rightarrow T(1, 1) = \mathbb{R}^4/\mathbb{Z}^4$ is injective on U .

The resulting embedding $B^4(a) \rightarrow T(1, 1)$ uses all four homological directions of $T(1, 1)$. This must be so. Indeed, assume that there exists an embedding $B^4(a) \rightarrow T(1, 1)$ that factors, for instance, as

$$B^4(a) \xrightarrow{\psi} T^3(x_1, y_1, x_2) \times (0, 1) \rightarrow T(1, 1)$$

with $(0, 1) \subset \mathbb{R}(y_2)$. It is easy to see that there exists a symplectic embedding ρ of the annulus $T^1(x_2) \times (0, 1)$ into $B^2(1) \subset \mathbb{R}^2(x_2, y_2)$. Composing ψ with $\text{id} \times \rho$ we obtain a symplectic embedding of $B^4(a)$ into $T^2(x_1, y_1) \times B^2(1)$, which lifts to $\mathbb{R}^2(x_1, y_1) \times B^2(1)$. Hence $a \leq 1$ by the Nonsqueezing Theorem.

Remark 1.8. Parts of the above construction yield an explicit full filling by one ball of the 4-torus $T(\mu, 1)$ for all $\mu = \frac{2m^2}{n^2}$ with m, n relatively prime. Since the set of rational numbers μ of this form is dense in $\mathbb{R}_{>0}$, one is tempted to derive $p(T(1, 1)) = 1$ from $p(T(\mu, 1)) = 1$ for $\mu > 1$ by a limiting argument, or to derive $p(T(\mu, 1)) = 1$ for all $\mu \geq 1$ from the elementary explicit full fillings of $T(\mu, 1)$ for $\mu = \frac{2m^2}{n^2}$. However, without further knowledge about the underlying embeddings, the function $\mu \mapsto p(T(\mu, 1))$ has no obvious continuity properties. $\diamond$

Filling by more than one ball. The general ball packing problem for a symplectic 4-manifold (M, ω) is: Given a collection $\overline{B}^4(a_1), \dots, \overline{B}^4(a_k)$ of closed balls, does there exist a symplectic embedding of $\coprod_{j=1}^k \overline{B}^4(a_j)$ into (M, ω) ? Since symplectic embeddings are volume preserving, a necessary condition is $\text{Vol}(\coprod_{j=1}^k \overline{B}^4(a_j)) < \text{Vol}(M, \omega)$. We prove that for all linear tori *except possibly* $T(1, 1)$ this is the only condition.

Theorem 1.9. *Assume that (T^4, ω) is a linear torus. Let $\overline{B}^4(a_1), \dots, \overline{B}^4(a_k)$ be a collection of balls such that*

$$\text{Vol}\left(\coprod_{j=1}^k \overline{B}^4(a_j)\right) < \text{Vol}(T^4, \omega).$$

- (i) *If (T^4, ω) is not symplectomorphic to $T(\mu, \mu)$ for some $\mu > 0$, there exists a symplectic embedding of $\coprod_{j=1}^k \overline{B}^4(a_j)$ into (T^4, ω) .*
- (ii) *If (T^4, ω) is symplectomorphic to $T(\mu, \mu)$ for some $\mu > 0$, then $\coprod_{j=1}^k \overline{B}^4(a_j)$ embeds symplectically into $T(\mu, \mu)$ under the further restriction that $a_j < \mu$ for all j .*

Notice that Theorem 1.9 generalizes Propositions 1.5 and 1.6. The extra condition in (ii) is presumably not needed, but the only way we can see to prove this would be by explicitly constructing suitable embeddings.

Other examples of manifolds for which the volume is the only obstruction to a symplectic embedding of a collection of balls were found by Biran in [4, 5]. Biran also proved in [5] that $T(1, 1)$ can be fully packed with k equal balls for any $k \geq 2$.³

Filling by ellipsoids. Recently, symplectic embeddings of ellipsoids into symplectic manifolds have attracted much attention, see [9, 10, 17, 35, 37, 40]. For $a, b > 0$ define the four-dimensional open symplectic ellipsoid⁴

$$E(a, b) = \left\{ (z_1, z_2) \in \mathbb{C}^2 \mid \frac{\pi|z_1|^2}{a} + \frac{\pi|z_2|^2}{b} < 1 \right\}.$$

The results in McDuff [35] apply to prove the following generalization of Theorem 1.9.

Theorem 1.10. *Assume that (T^4, ω) is a linear torus. Let $\overline{E}(a_1, b_1), \dots, \overline{E}(a_k, b_k)$ be a collection of ellipsoids such that*

$$\text{Vol} \left(\coprod_{j=1}^k \overline{E}(a_j, b_j) \right) < \text{Vol} (T^4, \omega).$$

- (i) *If (T^4, ω) is not symplectomorphic to $T(\mu, \mu)$ for some $\mu > 0$, there exists a symplectic embedding of $\coprod_{j=1}^k \overline{E}(a_j, b_j)$ into (T^4, ω) .*
- (ii) *If (T^4, ω) is symplectomorphic to $T(\mu, \mu)$ for some $\mu > 0$, then $\coprod_{j=1}^k \overline{E}(a_j, b_j)$ embeds symplectically into $T(\mu, \mu)$ under the further restriction that $\min\{a_j, b_j\} < \mu$ for all j .*

We shall prove Theorem 1.10 first for irrational tori, and then for product tori $T(\mu, 1)$ under the condition that $\min\{a_j, b_j\} < \mu$ for all j . This extra condition follows from the volume condition when $\mu \geq 2$. In view of Lemma 1.4 the theorem then follows. For embeddings of one ellipsoid, the proviso in (ii) means that we do not know whether $T(1, 1)$ can be filled by a dilate of $E(1, b)$, $b \geq 1$, if $b \in (1, 2)$. (For $b = 1$, we obtain a full filling by an explicit construction.)

Remark 1.11. Another important invariant of a symplectic manifold (M, ω) is its Hofer–Zehnder capacity $c_{\text{HZ}}(M, \omega)$, which is of dynamical nature. We refer to the books [16, 39] for the definition and elementary properties. The value of this capacity is unknown for product tori; in fact it is an outstanding problem to decide whether it is finite or infinite for product tori.

Our computations of the Gromov width c_G of tori give lower bounds for c_{HZ} , because $c_G(M, \omega) \leq c_{\text{HZ}}(M, \omega)$ for all symplectic manifolds. These lower bounds are, however, weaker than the known ones. These come from the elementary inequality

$$c_{\text{HZ}}(M, \omega) = c_{\text{HZ}}(P, \omega_P) + c_{\text{HZ}}(Q, \omega_Q),$$

³In fact, his argument also proves the claim in Theorem 1.9 concerning $T(1, 1)$. His proof is much the same as ours in that he reduces the problem to packing some ruled 4-manifold. However he considers the projectivization of a holomorphic line bundle of Chern class 2 over a genus 2 surface, while we use a trivial bundle over T^2 . In both cases the spherical fibers have area 1.

⁴More generally one might define a symplectic ellipsoid to be a set of the form $Q(x) < 1$, where Q is a positive definite quadratic form on $\mathbb{R}^4$. But this gives nothing new since every such Q can be diagonalized by a symplectic linear map.

holding for all products $(M, \omega) = (P \times Q, \omega_P \oplus \omega_Q)$ of closed symplectic manifolds, together with the fact that the Hofer–Zehnder capacity of a 2-dimensional connected symplectic manifold equals its area. To be explicit, our main theorem implies that $c_{\text{HZ}}(T(1, 1)) \geq c_G(T(1, 1)) = \sqrt{2}$, while it is known that $c_{\text{HZ}}(T(1, 1)) \geq 1 + 1 = 2$. $\diamond$

The paper is organized as follows. In Section 2 we review the lower bounds for the ball filling number of 4-dimensional symplectic tori coming from known computations of Seshadri constants. Section 3.1 contains proofs of Theorems 1.9 and 1.10 in the irrational case. These proofs are based on the construction in Section 3.2 of symplectic tori with no holomorphic curves. In Section 3.3 we prove Theorems 1.9 and 1.10 for product tori $T(1, \mu)$, $\mu \geq 1$, under the condition that $\min\{a_j, b_j\} < \mu$ for all j . In Section 4 we explain the embedding construction that we use in Section 5 to prove $p(T(1, 1)) = 1$, completing the proof of Theorem 1.1. In Section 6 we prove Corollaries 1.2 and 1.3, and in Section 7 we state some open problems related to filling tori.

Acknowledgment. This work has its origin in discussions between the authors and Dietmar Salamon at the Edifest 2010, and we would like to thank ETH Zürich and its FIM for the stimulating atmosphere during the conference. We also thank Paul Biran, Dietmar Salamon and Sewa Shevchishin for fruitful discussions.

2. RELATIONS TO ALGEBRAIC GEOMETRY

In this section, we review the implications of some results in algebraic geometry for the Gromov radius of 4-dimensional symplectic tori, and also of some higher dimensional ones.

2.1. Basics. Before discussing the complex geometry of tori, we recall a classical result.

Lemma 2.1. *Suppose ω is a linear symplectic form on a torus $T = \mathbb{R}^{2n}/\Lambda$ with integral cohomology class. Then (T, ω) is symplectomorphic to a product of 2-dimensional tori*

$$T^2(d_1) \times \cdots \times T^2(d_n)$$

with symplectic areas $d_j > 0$ satisfying $d_j | d_{j+1}$ for all $j = 1, \dots, n-1$. Moreover, the sequence $d_1 | d_2 | \dots | d_n$ is uniquely determined by Λ .

Remark 2.2. It follows that a 4-dimensional product torus $T(m, n) = T^2(m) \times T^2(n)$ with integer areas m and n is symplectomorphic to $T(g, \ell)$ where $g = \gcd(m, n)$, $\ell = \text{lcm}(m, n)$.

Proof of Lemma 2.1. Since ω is linear, it lifts to a linear symplectic form on $\mathbb{R}^{2n}$, which we again denote by ω . The fact that it represents an integral cohomology class on T is equivalent to the fact that it takes integer values when restricted to $\Lambda \times \Lambda$. Denote by $d_1 \in \mathbb{Z}$ the positive generator of this image subgroup, and choose $e_1, f_1 \in \Lambda$ with $\omega(e_1, f_1) = d_1$.

Every lattice point $\lambda \in \Lambda$ can be written as

$$\lambda = \frac{\omega(\lambda, f_1)}{d_1} e_1 + \frac{\omega(e_1, \lambda)}{d_1} f_1 + \lambda',$$

where the coefficients of e_1 and f_1 are integers by the choice of d_1 , and where $\lambda' \in \Lambda$ is ω -orthogonal to both e_1 and f_1 . In other words, $\Lambda = \text{span}_{\mathbb{Z}}(e_1, f_1) \oplus \Lambda'$ for some lower dimensional sublattice $\Lambda' \subset \Lambda$. Now repeat the argument with Λ' in place of Λ , noting that the image of ω when restricted to $\Lambda' \times \Lambda'$ must be a subgroup of $d_1 \mathbb{Z} \subset \mathbb{Z}$. This finishes the proof in n steps.

To prove the uniqueness of the sequence $d_1 | \dots | d_n$ for a given torus $T = \mathbb{R}^{2n}/\Lambda$, note that since ω is non-degenerate and integral, it gives rise to an embedding $\phi : \Lambda \rightarrow \text{Hom}(\Lambda, \mathbb{Z})$, namely $\phi(\lambda_1)(\lambda_2) = \omega(\lambda_1, \lambda_2)$. Now the d_j are the torsion coefficients of the finitely generated abelian group $\text{Hom}(\Lambda; \mathbb{Z})/\text{Im } \phi$, which are well-known to be invariants of this group. $\square$

Complex tori are often defined as the quotients of $\mathbb{C}^n$ by some cocompact lattice $\Lambda \cong \mathbb{Z}^{2n}$. In dimension 4, the Enriques–Kodaira classification of compact complex surfaces implies that every complex manifold diffeomorphic to T^4 is biholomorphic to such a model. In higher dimensions, this is still true if the complex structure is compatible with a Kähler form, but false in general (for examples, see e.g. [11] and references therein).

Conversely, the standard symplectic form on $\mathbb{C}^n$ descends to a Kähler form on any quotient $\mathbb{C}^n/\Lambda$, so every complex torus admits a compatible Kähler structure whose symplectic form is translation invariant.

2.2. Seshadri constants of tori. Here we review some results described by Lazarsfeld in [26, Chapter 5], which do not seem to be widely known among symplectic geometers. For an irreducible projective variety X and a point $x \in X$ we denote by

$$\pi : \tilde{X} \rightarrow X$$

the blow-up of X at x , with exceptional divisor $\Sigma \subset \tilde{X}$. Recall that a line bundle L on X is called *nef* if for every irreducible curve $C \subset X$ one has $\int_C c_1(L) \geq 0$.

Definition 2.3. (cf. [26, Def. 5.1.1.]) Suppose L is a nef line bundle on X . The *Seshadri constant of (X, L) at $x \in X$* is defined to be the real number

$$\varepsilon(L; x) := \max\{\varepsilon \geq 0 \mid \int_{\tilde{C}} \pi^*(c_1 L) - \varepsilon \Sigma \cdot \tilde{C} \geq 0 \text{ for all curves } \tilde{C} \subset \tilde{X}\}.$$

It is clear that $\varepsilon(L, x)$ is always nonnegative, and in fact one has the alternative description (cf. [26, Prop. 5.1.5.])

$$(2.1) \quad \varepsilon(L; x) = \inf_{x \in C \subset X} \frac{\int_C c_1(L)}{\text{mult}_x C},$$

where the infimum is taken over all irreducible curves $C \subset X$ passing through x , and $\text{mult}_x C \in \mathbb{N}$ denotes the multiplicity of C at x . This shows that one can obtain upper bounds on $\varepsilon(L; x)$ from specific curves passing through $x \in X$.

From the symplectic point of view, we are particularly interested in the case when X is a smooth projective variety, and L is an ample line bundle. Then one can choose a Kähler form ω_L representing $c_1(L)$. Since the space of symplectic forms in a fixed cohomology class which are compatible with a fixed (almost) complex structure is contractible, any two such forms are symplectically isotopic.

Now there is a strong relationship between symplectic embeddings of balls and symplectic blow-up, which was first described by McDuff [30] and McDuff–Polterovich [38]. Basically, an embedding of a closed symplectic ball $B(a)$ of capacity a into a given symplectic manifold X gives rise to a symplectic form on the blow-up $\pi : \tilde{X} \rightarrow X$ whose cohomology class is given by $\pi^*[\omega] - a \text{PD}[\Sigma]$, where $\Sigma \subset \tilde{X}$ is the exceptional divisor, and $\text{PD}[\Sigma]$ denotes the Poincaré dual of $[\Sigma]$. Conversely, given a symplectic form on $\tilde{X}$ in a class $\pi^*\alpha - a \text{PD}[\Sigma]$, one can find a symplectically embedded ball $B(a)$ in (X, ω) with $[\omega] = \alpha \in H^2(X; \mathbb{R})$.

It is important to observe that if one starts from a Kähler form on $\tilde{X}$, then the resulting form on X may also be constructed to be Kähler. As pointed out in [26, Thm. 5.1.22.], this discussion then leads to the following result, which is a direct consequence of [38, Cor. 2.1.D]:

Proposition 2.4. *For fixed X and L as above, denote by*

$$\varepsilon(X, L) := \max_{x \in X} \varepsilon(L; x).$$

Then the Gromov width of (X, ω_L) , defined in (1.1), satisfies

$$c_G(X, \omega_L) \geq \varepsilon(X, L).$$

By the relation (1.2), this estimate is equivalent to

$$(2.2) \quad p(X, \omega_L) \geq \frac{(\varepsilon(X, L))^n}{n! \operatorname{Vol}(X, \omega_L)}.$$

The proof of Proposition 2.4 is based on the fact that when $\varepsilon(X; L) > 0$, then the pullback $\tilde{L}$ of L to the blow-up $\tilde{X}$ is ample, so that $c_1(\tilde{L})$ has a Kähler representative.

Remark 2.5. The same blow-up argument also shows that the capacity of the largest *symplectically and holomorphically* embedded ball in the Kähler manifold (X, ω_L) bounds the Seshadri constant $\varepsilon(X, L)$ from below (for details, cf. [26, Prop. 5.3.17]).

In what follows, we will study the family of symplectic product tori $T^2(1) \times T^2(d)$ where $d \in \mathbb{Z}$. By Lemma 2.1, up to rescaling this class contains all symplectic 4-tori whose symplectic form is linear and has a rational cohomology class. Now suppose that (T, ω) is such a symplectic torus, and choose a translation-invariant compatible complex structure J , so that (T, J, ω) is a Kähler manifold. If L is the complex line bundle on (T, J) with first Chern class $[\omega]$, then L is ample. Complex tori admitting such a line bundle are called abelian varieties, and the line bundle or its first Chern class is called a *polarization*. Note that, conversely, the first Chern class of any ample line bundle L on some complex torus can be represented by a translation-invariant rational symplectic form ω_L , and so all abelian varieties arise as above.

The sequence of integers $(d_1, \dots, d_n)$ for (T, ω_L) appearing in Lemma 2.1 is called the *type* of the polarization, and the polarization is called *principal* and often denoted by Θ if it is of type $(1, \dots, 1)$, i.e. it corresponds to the standard symplectic product torus.

Since translations act transitively on any abelian variety A , the Seshadri constants for abelian varieties do not depend on the choice of the point $x \in A$. One has the general bounds (cf. [26, Prop. 5.1.9 and Ex. 5.3.10])

$$(2.3) \quad d_1 \leq \varepsilon(A^n, L) \leq (n! d_1 \cdots d_n)^{\frac{1}{n}}$$

for an ample line bundle L of type $(d_1, \dots, d_n)$. Note that the upper bound follows from the estimate (2.2). Similarly, the symplectic embedding of the ball of capacity d_1 into the polydisk $B^2(d_1) \times \cdots \times B^2(d_n) \subset A$ gives the same lower bound d_1 for the Gromov width of (A, ω_L) .

The best lower bounds on Seshadri constants for abelian varieties of a given type seem to come from irreducible ones, i.e., those which cannot be written as a product of lower-dimensional complex tori. Here we list the known results, according to [26, Rem. 5.3.12].

First, to get a bound on the ball filling number of $T(1, 1)$, according to the discussion above we need to consider principally polarized abelian surfaces (A^2, Θ) . Steffens [43, Prop. 2 and 3]

has shown that in this case

$$(2.4) \quad \varepsilon(A^2, \Theta) \leq \frac{4}{3},$$

with equality if A is irreducible. Together with the estimate (2.2) we obtain the lower bound

$$(2.5) \quad p(T(1, 1)) \geq \frac{8}{9}.$$

For tori of type $(1, d)$ one can get lower bounds from non-principal polarizations of abelian surfaces (A^2, L) . Indeed, it is known from the work of Steffens [43, Prop. 1] that *if $2d$ is a perfect square*, then there are abelian surfaces with a polarization L of type $(1, d)$ and

$$(2.6) \quad \varepsilon(A^2, L) = \sqrt{2d},$$

which is optimal since it equals the volume bound in (2.3). This immediately implies

$$(2.7) \quad p(T(1, d)) = 1 \quad \text{if } 2d \text{ is a perfect square.}$$

We will describe explicit examples of such full fillings by one symplectic ball in Section 4.3. The identities (2.7) and Remark 2.2 imply that $p(T(\mu, 1)) = 1$ for all $\mu = \frac{2m^2}{n^2}$ with m, n relatively prime integers.

On the other hand, when $2d$ is not a perfect square, then Bauer and Szemberg [1] have shown that

$$(2.8) \quad \varepsilon(A^2, L) \leq 2d \frac{k_0}{\ell_0} = \sqrt{2d} \cdot \sqrt{\frac{2dk_0^2}{2dk_0^2 + 1}},$$

where (k_0, ℓ_0) is the smallest solution in positive integers of Pell's equation

$$\ell^2 - 2dk^2 = 1.$$

(There always exists such a solution, as was first shown by Lagrange, [20].) Moreover, by a result of Bauer [2] equality holds whenever positive multiples of L are the only ample line bundles on A . Since complex structures J with this property exist for all symplectic types $(1, d)$, this gives the best constant for use in Proposition 2.4. For $d \leq 30$, the relevant solutions to Pell's equation have been tabulated in [2, p. 572], and we give their translation in terms of the lower bound on the ball filling numbers $p(T(1, d)) \geq \frac{\varepsilon^2}{2d} = \frac{\ell_0^2 - 1}{\ell_0^2}$ in the following table.

d	k_0	ℓ_0	$\frac{\varepsilon^2}{2d}$	d	k_0	ℓ_0	$\frac{\varepsilon^2}{2d}$	d	k_0	ℓ_0	$\frac{\varepsilon^2}{2d}$
1	2	3	$\frac{8}{9}$	11	42	197	$\frac{38808}{38809}$	21	2	13	$\frac{168}{169}$
2			1	12	1	5	$\frac{24}{25}$	22	30	199	$\frac{39600}{39601}$
3	2	5	$\frac{24}{25}$	13	10	51	$\frac{2600}{2601}$	23	3588	24335	$\frac{592192224}{592192225}$
4	1	3	$\frac{8}{9}$	14	24	127	$\frac{16128}{16129}$	24	1	7	$\frac{48}{49}$
5	6	19	$\frac{360}{361}$	15	2	11	$\frac{120}{121}$	25	14	99	$\frac{9800}{9801}$
6	2	7	$\frac{48}{49}$	16	3	17	$\frac{288}{289}$	26	90	649	$\frac{421200}{421201}$
7	4	15	$\frac{224}{225}$	17	6	35	$\frac{1224}{1225}$	27	66	485	$\frac{235224}{235225}$
8			1	18			1	28	2	15	$\frac{224}{225}$
9	4	17	$\frac{288}{289}$	19	6	37	$\frac{1368}{1369}$	29	2574	19605	$\frac{384356024}{384356025}$
10	2	9	$\frac{80}{81}$	20	3	19	$\frac{360}{361}$	30	4	31	$\frac{960}{961}$

2.3. Seshadri estimates for higher dimensional tori. One well-studied class of principally polarized abelian varieties of arbitrary dimension are the Jacobians of curves (cf. e.g. [7, Chapter 11]). Here we just recall that the Jacobian of a complex curve C is the complex torus

$$JC := \text{Hom}(\Omega^{1,0}, \mathbb{C})/H_1(C; \mathbb{Z}),$$

where $\Omega^{1,0}$ denotes the complex vector space of holomorphic 1-forms, and the embedding $H_1(C; \mathbb{Z}) \subset \text{Hom}(\Omega^{1,0}, \mathbb{C})$ is given by integration over cycles. The complex dimension of JC equals the genus of C , and the principal polarization is derived from the natural symplectic structure on $H_1(C; \mathbb{Z}) \otimes \mathbb{R}$ which is given by the intersection product.

In complex dimension $n = 3$, Bauer and Szemberg [3] have shown that a principally polarized abelian variety (A^3, Θ) has $\varepsilon(\Theta) = \frac{3}{2}$ if A is the Jacobian of a hyperelliptic curve of genus 3 and

$$(2.9) \quad \varepsilon(A^3, \Theta) = \frac{12}{7}$$

otherwise. (A complex curve is called hyperelliptic if it admits a double branched cover to CP^1). Hence $p(T(1, 1, 1)) \geq \frac{288}{343}$.

In complex dimension $n = 4$, Debarre [12] has shown that for the Jacobian $A^4 = JC$ of a non-hyperelliptic curve of genus 4 one has

$$(2.10) \quad \varepsilon(A^4, \Theta) = 2.$$

Hence $p(T(1, 1, 1, 1)) \geq \frac{2}{3}$.

For high dimensions, Jacobians appear to give very poor lower bounds for use in Proposition 2.4. However, Lazarsfeld [25] combined the work of McDuff and Polterovich [38] with work of Buser and Sarnak on minimal period lengths to deduce that there exist principally polarized abelian varieties (A^n, Θ) of complex dimension n with

$$\varepsilon(A^n, \Theta) \geq \frac{1}{4}(2n!)^{\frac{1}{n}}.$$

Bauer has generalized this, showing that there exist polarized abelian varieties (A^n, L) of arbitrary type $(d_1, \dots, d_n)$ with

$$(2.11) \quad \varepsilon(A^n, L) \geq \frac{1}{4}(2n!d_1 \dots d_n)^{\frac{1}{n}}.$$

While this is only a factor of less than 4 away from the upper bound of (2.3), the volume fraction filled by the symplectic ball predicted from this lower bound is $2\left(\frac{1}{4}\right)^n$, and hence tends exponentially to zero as $n \rightarrow \infty$.

3. PROOF OF THEOREMS 1.9 AND 1.10

3.1. Irrational case. We will use a result of Buchdahl [8] and Lamari [24], which states that a cohomology class $\alpha \in H^{1,1}(X; \mathbb{R})$ on some complex surface X with even first Betti number admits a Kähler representative if $\alpha \cup \alpha > 0$, $\alpha \cup [\rho] > 0$ for some positive closed $(1,1)$ -form ρ on X and $\alpha \cdot [D] > 0$ for every effective divisor $D \subset X$. In symplectic language, the last condition means that the class α should integrate positively over every compact holomorphic curve in X .

Our argument is based on the following result, whose proof is deferred until the next subsection.

Proposition 3.1. *Any irrational linear symplectic form ω on T^4 may be identified with a Kähler form on a torus $T = \mathbb{C}^2/\Lambda$ that has no nonconstant compact holomorphic curves.*

Proposition 3.2. *Theorem 1.9 holds for irrational tori.*

Proof. We must show that any disjoint union $\coprod_{j=1}^k \overline{B}^4(a_j)$ of balls embeds symplectically into (T^4, ω) provided only that the volume constraint is satisfied. By Proposition 3.1 there is a symplectomorphism from (T^4, ω) to (T, ω_J) where (T, J, ω_J) is the Kähler torus found in Proposition 3.1. Let (T_k, J_k) be the complex blow-up of T at k generic points, and consider the cohomology class $\alpha := \pi^*([\omega_J]) - \sum_{j=1}^k a_j \text{PD}[\Sigma_j]$, where $\pi: T_k \rightarrow T$ is the blow-down map and the Σ_j are the exceptional divisors. Choose $\varepsilon > 0$ so small that there exist disjoint symplectically and holomorphically embedded balls $B^4(\varepsilon)$ around the k blow-up points, and that $\int_T \omega_J \cup \omega_J > \sum_{j=1}^k \varepsilon a_j$. Let ρ be the Kähler form on T_k corresponding to the blow-up of these k balls. Then $[\rho] = \pi^*([\omega_J]) - \sum_{j=1}^k \varepsilon \text{PD}[\Sigma_j]$, whence $\alpha \cup [\rho] > 0$. Furthermore, the volume condition gives $\alpha^2 > 0$, and the only complex curves in (T_k, J_k) are the Σ_j . The criterion of Buchdahl and Lamari thus holds for the class α . Therefore there is a Kähler form on (T_k, J_k) in the class α .

Now blow down this form⁵ to obtain a Kähler form η on (T, J) in the class $[\omega_J]$, such that (T, η) contains disjoint holomorphically and symplectically embedded balls $\overline{B}^4(a_1), \dots, \overline{B}^4(a_k)$. Averaging over translations if necessary, we obtain a linear symplectic form η' in the same cohomology class. Hence $\eta' = \omega_J$. Further, since η' is compatible with the complex structure of T , we can linearly interpolate η and η' by symplectic forms, and use Moser's argument to see that (T, η) and (T, ω_J) are symplectomorphic. Hence the balls $\coprod_j \overline{B}^4(a_j)$ also embed symplectically in (T, ω_J) , but perhaps not holomorphically. $\square$

Proposition 3.3. *Theorem 1.10 holds for irrational tori.*

Proof. This is very similar to the proof of Theorem 1.9 except for the last step. For simplicity, we explain the proof for one ellipsoid. The proof for more ellipsoids is essentially the same.

First note that by a simple continuity argument (as in [40, Lemma 1.1]) it suffices to consider the case when the ellipsoid $\overline{E}(a, b)$ has rational ratio a/b . It is explained in [35] how to convert an ellipsoidal embedding problem into a question about constructing a suitable symplectic form on a blow-up: an embedded ellipsoid $\lambda \overline{E}(m, n) := \overline{E}(\lambda m, \lambda n)$ with $m, n \in \mathbb{N}$ corresponds

⁵The Kähler blow-down is discussed for example in [26, Prop. 5.3.15].

to a pair consisting of a singular reducible curve $C(m, n)$ in the blow-up, whose properties are determined by the + continued fraction expansion of m/n , together with a symplectic form on the blow-up that is nondegenerate on $C(m, n)$ and lies in a suitable cohomology class. For example the ellipsoid $\lambda\overline{E}(1, 3)$ corresponds to a chain of spheres in classes $E_1 - E_2, E_2 - E_3, E_3$, where each E_i is the class of an exceptional divisor, and the corresponding form has integral just more than λ on E_3 and is very small on the other components. In all cases, $C(m, n)$ is a chain of pairwise intersecting embedded 2-spheres, and can be created by suitable nongeneric complex blow-up (in fact one is just resolving certain toric singularities; see [36, §2.2]). Hence, just as in the previous proof one can use the Buchdahl–Lamari criterion to construct a suitable Kähler form τ on the blow-up.

The new point comes in the blow-down construction. So far, only a symplectic version of this construction has been worked out. The idea is to perturb the components of the configuration $C(m, n)$ so that the components of the resulting configuration $C'(m, n)$ intersect orthogonally with respect to the Kähler form τ . Then a neighborhood of $C'(m, n)$ is symplectomorphic to a standard toric model, and so it can be blown down using the so-called “rational blow-down”; cf. [35, Lemma 2.3]. The difficulty now is to prove that the resulting form on the blow-down is symplectomorphic to a linear form. One way to deal with this is to perform this construction for the family of ellipsoids $s\overline{E}(a, b)$, $0 < s \leq 1$. Correspondingly we have a family of Kähler forms τ_s , $0 < s \leq 1$, on the blow-up. (Since the set of Kähler forms in a given cohomology class is convex, it is easy to see that we can choose the τ_s to vary smoothly with $s > 0$.) We can find deformations of $C(m, n)$ to $C'_s(m, n)$ that depend smoothly on s for $s > 0$ and then get a smooth family of cohomologous symplectic forms η_s , $s > 0$, on the blow-down. By construction, the ellipsoid $s\overline{E}(a, b)$ embeds symplectically into (T, η_s) for all $s \in (0, 1]$. The only problem is to guarantee that these η_s converge smoothly as $s \rightarrow 0$ to some form η_0 on T that is isotopic (and hence symplectomorphic) to the original linear form ω_J . For if this were true, η_1 would be isotopic to ω_J , so that there would be a symplectic embedding of $\overline{E}(a, b)$ into (T, ω_J) .

We now show that ω_J can be isotoped to a J -tame form ω'_J such that for some $\varepsilon > 0$ we have $\eta_s = \omega'_J$ for $s \in (0, \varepsilon]$. First of all, given any point $x \in T$ we can choose a local chart $\psi: (U_x, x) \rightarrow (U_0, 0)$, where U_x is a neighborhood of x in T and U_0 is a neighborhood of $0 \in \mathbb{C}^2$, which is holomorphic, i.e. $\psi_*(J_T) = J_0$, and is such that $\psi_*(\omega_T) =: \omega$ equals the standard form ω_{can} at the point 0. (This follows from [39, Exercise 2.52 (iii)].) Then a standard Moser argument allows us to find a family of symplectic forms ω_t , $t \in [0, 1]$, on U_0 such that

- $\omega_0 = \omega$ and $\omega_t = \omega$ near ∂U_0 ;
- ω_t is J_0 -tame for all t ;
- there is $\delta > 0$ such that $\omega_1 = \omega_{\text{can}}$ in $B^4(\delta) \subset U_0$.

Then for any $s > 0$ such that $s\overline{E}(a, b) \subset B^4(\delta)$ we can blow up symplectically at x to get the configuration $C(m, n)$ with appropriate Kähler form, and then blow down in such a way that the blow-down form on U is the one we started with. Hence if we take $\omega'_J = \psi^*(\omega_1)$ we may choose the path τ_s so that for some short interval $s \in (0, \varepsilon]$ it consists of forms that blow down to ω'_J . This completes the proof. $\square$

3.2. Irrational tori with no curves.

We now prove Proposition 3.1.

We begin by finding complex tori with no nonconstant compact holomorphic curves. Consider $\mathbb{C}^2 = \mathbb{C}e_1 \oplus \mathbb{C}e_2$ and denote by $e_1, e_2, e_3 = \sqrt{-1}e_1, e_4 = \sqrt{-1}e_2$ the standard real basis.

Choose real numbers p, q, r, s such that

$$(3.1) \quad p, q, r, s \text{ are rationally independent, and } ps - qr \text{ is irrational.}$$

Consider the quotient of $\mathbb{C}^2$ by the lattice Λ_P spanned by

$$\lambda_1 := e_1, \quad \lambda_2 := e_2, \quad \lambda_3 = p e_3 + q e_4, \quad \lambda_4 = r e_3 + s e_4.$$

The following result is taken from the Appendix in [14]. We repeat it here for the sake of completeness.

Lemma 3.4. *Under the assumption (3.1), the torus $T = \mathbb{C}^2/\Lambda_P$ contains no nonconstant compact holomorphic curves.*

Proof. Any such curve would represent a class in $H_{1,1}(T; \mathbb{C}) \cap H_2(T; \mathbb{Z})$. By duality it suffices to prove that $H^2(T; \mathbb{Z}) \cap H^{1,1}(T; \mathbb{C}) = \{0\}$. By the Lefschetz Theorem on $(1, 1)$ -classes this is the Néron–Severi group of T . It can be identified with the set of Hermitian forms H on $\mathbb{C}^2$ satisfying $\text{im } H(\Lambda_P \times \Lambda_P) \subset \mathbb{Z}$. In our standard real basis of $\mathbb{C}^2$, the period matrix of Λ_P is

$$P = \begin{pmatrix} 1 & 0 & \sqrt{-1}p & \sqrt{-1}r \\ 0 & 1 & \sqrt{-1}q & \sqrt{-1}s \end{pmatrix},$$

and each Hermitian form is given by a matrix

$$(3.2) \quad A = \begin{pmatrix} x & u + \sqrt{-1}v \\ u - \sqrt{-1}v & y \end{pmatrix}$$

with x, y, u, v real. The condition $\text{im } H(\Lambda_P \times \Lambda_P) \subset \mathbb{Z}$ becomes $\text{im } P^T A \overline{P} \in \mathbb{Z}^4 \times \mathbb{Z}^4$. Since

$$(3.3) \quad \text{im } P^T A \overline{P} = \begin{pmatrix} 0 & v & -(px + qu) & -(tx + su) \\ -v & 0 & -(pu + qy) & -(ru + sy) \\ px + qu & pu + qy & 0 & v(ps - qr) \\ rx + su & ru + sy & -v(ps - qr) & 0 \end{pmatrix},$$

this integrality condition is

- (i) $v, v(ps - qr) \in \mathbb{Z};$
- (ii) $px + qu, rx + su, pu + qy, ru + sy \in \mathbb{Z}.$

Since we have chosen $ps - qr$ irrational, the two conditions (i) imply $v = 0$. Assume that x, y, u fulfill the conditions (ii). We then find $n_1, n_2, n_3, n_4 \in \mathbb{Z}$ with

$$\begin{aligned} px + qu &= n_1, & pu + qy &= n_3, \\ rx + su &= n_2, & ru + sy &= n_4. \end{aligned}$$

We can eliminate x and y from the above equations and obtain

$$\begin{aligned} (ps - qr)u &= -n_1r + n_2p \\ (ps - qr)u &= n_3s - n_4q, \end{aligned}$$

which implies $-n_1r + n_2p - n_3s + n_4q = 0$. Since we have chosen p, q, r, s to be rationally independent, it follows that n_1, n_2, n_3, n_4 must vanish. Hence u and therefore also x and y vanish. We conclude that $A = 0$, as we wanted to show. $\square$

Now we start with a torus $T^4 = \mathbb{R}^4/\Lambda$ with a linear symplectic form ω representing an irrational cohomology class. Given an integral basis $\lambda_1, \dots, \lambda_4$ for Λ , the symplectic form ω can be represented by a matrix $B = (b_{ij})$, where $b_{ij} = \omega(\lambda_i, \lambda_j)$. We denote by $\lambda_1^*, \dots, \lambda_4^*$ the basis dual to $\lambda_1, \dots, \lambda_4$, and we may assume that the ordering has been chosen such that $\omega \wedge \omega$ is a positive multiple of $\lambda_1^* \wedge \lambda_3^* \wedge \lambda_2^* \wedge \lambda_4^*$.

Lemma 3.5. *In the situation just described, after changing the basis of $\mathbb{R}^4$ by an element of $\mathrm{SL}(4, \mathbb{Z})$, we may represent ω by a matrix B' where*

- (i) *the entries b'_{12}, b'_{34} either both vanish or they are rationally independent and positive, and*
- (ii) *the vector $(b'_{13}, b'_{14}, b'_{23}, b'_{24})$ is not a multiple of a rational vector.*

Proof. Suppose first that there is a permutation $i_1, \dots, i_4$ of $\{1, \dots, 4\}$ so that $b_{i_1 i_2} = b_{i_3 i_4} = 0$. Then we can change basis (preserving orientation) so that $b'_{12} = b'_{34} = 0$. Condition (ii) is then automatic since ω is irrational.

In all other cases we can permute the basis (preserving orientation) so that $b_{12} \neq 0, b_{34} \neq 0$, and so that the elements b_{12}, b_{34}, b_{1j} are rationally independent for $j = 3$ or 4 . If condition (i) is not satisfied, we change basis by replacing λ_2^* by $\lambda_2^* + k\lambda_j^*$ and leaving the other elements fixed. Then b_{12} changes to $b'_{12} = b_{12} + kb_{1j}$ and $b'_{34} = b_{34}$, so that for suitable $k \in \mathbb{Z}$ we may assume that b'_{12}, b'_{34} are rationally independent and of the same sign. If they are both negative, we can change their signs by interchanging λ_1 and λ_2 and interchanging λ_3 and λ_4 . This achieves (i).

If (ii) does not hold, we may assume that the vector $\beta = (b_{13}, b_{14}, b_{23}, b_{24})$ does not vanish, because otherwise by a permutation we could have arranged the situation with $b'_{12} = b'_{34} = 0$ from the beginning of the proof, and as observed there (ii) is automatic in that case.

So one of the entries of β must be nonzero and hence rationally independent of either b_{12} or b_{34} . We will consider the case that b_{13} and b_{12} are rationally independent, the other cases being treated in a similar fashion. Now we change basis, replacing λ_4^* by $\lambda_4^* + k\lambda_1^*$ and leaving the other elements fixed. Then $b'_{ij} = b_{ij}$ if $i, j \neq 4$, while $b'_{i4} = b_{i4} - kb_{i1}$. In particular,

$$b'_{12} = b_{12}, \quad b'_{34} = b_{34} - kb_{13}, \quad b'_{13} = b_{13}, \quad b'_{24} = b_{24} - kb_{12}.$$

Hence (ii) holds if $k \neq 0$, since b'_{13}, b'_{24} are rationally independent. Further (i) will hold if we choose k so that $-kb_{13} > 0$.

The proof in the other cases is similar. In particular if $b_{13} = 0$ but $b_{14} \neq 0$ we use a base change that alters λ_3^* instead of λ_4^* . $\square$

Remark 3.6. Note that

$$\omega \wedge \omega = (b_{13}b_{24} - b_{14}b_{23} - b_{12}b_{34}) \lambda_1^* \wedge \lambda_3^* \wedge \lambda_2^* \wedge \lambda_4^*.$$

Since the base change was orientation preserving, the coefficient is still positive, and so in the new basis for Λ constructed in Lemma 3.5 we necessarily have $b'_{13}b'_{24} - b'_{14}b'_{23} > 0$, since $-b'_{12}b'_{34} \leq 0$ by (i).

Proof of Proposition 3.1. We are given a torus $T^4 = \mathbb{R}^4/\Lambda$ with a linear irrational symplectic form. We assume that we have chosen a basis $\lambda_1, \dots, \lambda_4$ for Λ such that the matrix B determined from $\omega(\lambda_i, \lambda_j) = b_{ij}$ satisfies the conditions stated in Lemma 3.5. Our goal is to identify Λ with a suitable period lattice Λ_P of the form discussed in Lemma 3.4, where the coefficients p, q, r, s are still to be determined, and to find a positive definite hermitian matrix A as in (3.2) above such that $B = -\mathrm{im}(P^T A \overline{P})$. With respect to the real standard basis

$e_1, e_2, e_3 = \sqrt{-1} e_1, e_4 = \sqrt{-1} e_2$ of $\mathbb{C}^2$ the metric g_A associated to the hermitian matrix A is represented by the matrix

$$(g_{ij}) = \begin{pmatrix} x & u & 0 & v \\ u & y & -v & 0 \\ 0 & -v & x & u \\ v & 0 & u & y \end{pmatrix}.$$

This matrix is positive definite if all leading principal minors are positive. This will be the case if and only if

$$(3.4) \quad x > 0 \quad \text{and} \quad xy - u^2 - v^2 > 0,$$

since the other two conditions $xy - u^2 > 0$ and $(xy - u^2 - v^2)^2$ are then necessarily also satisfied.

To identify B with $-\text{im}(P^T A \bar{P})$, the computation in equation (3.3) of the proof of Lemma 3.4 shows that we want

$$\begin{aligned} b_{12} &= -v, & b_{13} &= px + qu, \\ b_{14} &= rx + su, & b_{23} &= pu + qy, \\ b_{24} &= ru + sy, & b_{34} &= -v(ps - qr). \end{aligned}$$

The middle four equations can be rewritten as

$$(3.5) \quad \begin{pmatrix} q & p & 0 \\ s & r & 0 \\ p & 0 & q \\ r & 0 & s \end{pmatrix} \begin{pmatrix} u \\ x \\ y \end{pmatrix} = \begin{pmatrix} b_{13} \\ b_{14} \\ b_{23} \\ b_{24} \end{pmatrix}.$$

Here the vector on the right hand side is given and nonzero. For fixed p, q, r and s this overdetermined system of equations will have a solution (u, x, y) if the compatibility condition

$$(3.6) \quad r b_{13} - p b_{14} = q b_{24} - s b_{23}$$

is satisfied.

Lemma 3.7. *Since the vector $(b_{13}, b_{14}, b_{23}, b_{24})$ is not a multiple of a rational vector, there exists a solution $(p', q', r', s') \in \mathbb{R}^4$ of (3.6) in rationally independent real numbers satisfying $s'b_{13} - q'b_{14} > 0$ and $D := p's' - q'r' > 0$.*

Proof. The inequalities $sb_{13} - qb_{14} > 0$ and $D > 0$ define an open set in $\mathbb{R}^4$, which we denote by $\mathcal{O}$. Similarly, for given b_{ij} the equation (3.6) defines a hyperplane H in $\mathbb{R}^4$. The intersection $\mathcal{O} \cap H$ is not empty, since in view of Remark 3.6 the point with coordinates $p = b_{13}$, $q = b_{23}$, $r = b_{14}$ and $s = b_{24}$ belongs to it.

On the other hand, a point (p, q, r, s) has rationally dependent coordinates if and only if it solves some equation $n_1 p + n_2 q + n_3 r + n_4 s = 0$ with integral coefficients n_i . Since its defining vector is not a (multiple of a) rational vector, H is transverse to this countable set of hyperplanes, and so there is some point in the open subset $\mathcal{O} \cap H$ that does not lie on any of these hyperplanes. This point has the desired properties. $\square$

Given (p', q', r', s') as in the lemma, the solution of the matrix equation (3.5) is

$$\begin{aligned} x' &= \frac{1}{D}(s'b_{13} - q'b_{14}), & y' &= \frac{1}{D}(p'b_{24} - r'b_{23}), \\ u' &= \frac{1}{D}(p'b_{14} - r'b_{13}) = \frac{1}{D}(s'b_{23} - q'b_{24}). \end{aligned}$$

If $b_{12} = b_{34} = 0$, the final two equations

$$b_{12} = -v, \quad b_{34} = -v(p's' - q'r'),$$

have the trivial solution $v = 0$. In this case choose $\rho > 0$ such that $\rho^2(p'q' - r's')$ is irrational, and define $(p, q, r, s) := \rho(p', q', r', s')$ and $(u, x, y) := \rho^{-1}(u', x', y')$.

If b_{12}, b_{34} are rationally independent, we need to rescale the above solution so that $ps - qr = \frac{b_{34}}{b_{12}}$. Therefore, define $(p, q, r, s) := \rho(p', q', r', s')$ and $(u, x, y) := \rho^{-1}(u', x', y')$, where $\rho := \sqrt{\frac{b_{34}}{b_{12}D}}$. Notice that $ps - qr = \frac{b_{34}}{b_{12}}$ is irrational by part (i) of Lemma 3.5 in this case. Now, choosing $v = -b_{12}$, all six equations are satisfied.

It follows that in both cases, $\omega = -\text{im } Q_A$ for the Hermitian form Q_A determined by A on $\mathbb{C}^2/\Lambda_P$. To check the positivity condition (3.4), note that $x > 0$ by construction, and a computation shows that $xy - u^2 - v^2$ equals a positive multiple of $b_{13}b_{24} - b_{14}b_{23} - b_{12}b_{34}$, which was observed to be positive in Remark 3.6. In summary, we have shown that ω is compatible with the standard complex structure J_0 on $\mathbb{C}^2$.

Finally, by Lemma 3.7 and the subsequent construction, the period matrix P of the lattice satisfies the assumption of Lemma 3.4, and so we have proven the proposition. $\square$

3.3. Product tori. Theorems 1.9 and 1.10 for the product tori $T(1, \mu)$ with $\mu \geq 1$ are an immediate consequence of the following proposition.

Proposition 3.8. *Let $\overline{E}(a_1, b_1), \dots, \overline{E}(a_k, b_k)$ be a collection of ellipsoids such that*

$$\text{Vol} \left(\coprod_{j=1}^k \overline{E}(a_j, b_j) \right) < \mu$$

and such that $\min\{a_j, b_j\} < \mu$ for all j . Then there exists a symplectic embedding of $\coprod_{j=1}^k \overline{E}(a_j, b_j)$ into the open disc bundle $T^2(1) \times D^2(\mu)$.

Proof. The idea of the proof is as follows: Denote by $S^2(\mu)$ the 2-sphere endowed with an area form of area μ . Biran has shown in [4, Proof of Corollary 5.C] that $\coprod_{j=1}^k \overline{B}^4(a_j)$ symplectically embeds into $T^2(1) \times S^2(\mu)$ whenever $\text{Vol} \left(\coprod_{j=1}^k \overline{B}^4(a_j) \right) < \mu$ and $a_j < \mu$ for all j . We will arrange this embedding in such a way that the balls lie in $T^2(1) \times (S^2(\mu) \setminus z_0) = T^2(1) \times D^2(\mu)$, where $z_0 \in S^2$. Furthermore, Biran's arguments also apply to ellipsoids since, as shown in [35], the question of whether one can embed a given family of ellipsoids into a ruled manifold is equivalent to the question of whether one can embed an associated family of balls, where the balls corresponding to $E(a_j, b_j)$ have size not greater than $\min\{a_j, b_j\}$.

We obtain a symplectic embedding of $\coprod_{j=1}^k \overline{E}(a_j, b_j)$ into $T^2(1) \times (S^2(\mu) \setminus z_0)$, by constructing a smooth family of cohomologous forms ω_s , $s \in [0, 1]$, on $T^2 \times S^2$ with the following properties:

- ω_0 is the product form on $T^2(1) \times S^2(\mu)$;
- each ω_s is nondegenerate on the torus $C := T^2 \times z_0$;

- for each $s \in [0, 1]$ there is a symplectic embedding of $\coprod_{j=1}^k s\overline{E}(a_j, b_j)$ into $(T^2 \times S^2) \setminus C$.

For then a standard Moser argument shows that there is a family of diffeomorphisms $\psi_s : (T^2 \times S^2, C) \rightarrow (T^2 \times S^2, C)$ such that $\psi_1^* \omega_1 = \omega_0$. Therefore $\coprod_{j=1}^k \overline{E}(a_j, b_j)$ embeds symplectically into $(T^2 \times (S^2 \setminus z_0), \omega_0) = T^2(1) \times D^2(\mu)$.

We construct the family ω_s in much the same way as in the proof of Theorem 1.10 for irrational tori. In other words, we convert the problem into one of constructing suitable forms τ_s on the appropriate blow-up. The only difference is that we can no longer find the required forms τ_s on the blow-up via the Buchdahl–Lamari criterion; instead we must use symplectic inflation as in [4, 33, 35]. In order for C to be τ_s -symplectic, it suffices to work only with almost complex structures J for which C is J -holomorphic. Since such methods have been used many times (for example in [5]), we leave further details to the reader. $\square$

4. BASIC SYMPLECTIC MAPPINGS

In this section we describe an elementary symplectic embedding construction. It will be applied in Section 5 to prove that $p(T(1, 1)) = 1$.

4.1. Diamonds. Consider the “diamond” of size a

$$\diamond(a) := \{(x_1, x_2) \in \mathbb{R}^2 \mid |x_1| + |x_2| < \frac{a}{2}\} \subset \mathbb{R}^2(x),$$

see Figure 3 (I).

Lemma 4.1. *For each $\varepsilon > 0$ the ball $B^4(a)$ symplectically embeds into $\diamond(a + \varepsilon) \times (0, 1)^2 \subset \mathbb{R}^2(x) \times \mathbb{R}^2(y)$.*

Proof. Let $D(a) \subset \mathbb{R}^2(z) = \mathbb{R}^2(x, y)$ be the open disc of area a . Choose an area and orientation preserving embedding

$$\sigma : D(a) \rightarrow \left(-\frac{a + \varepsilon}{2}, \frac{a + \varepsilon}{2}\right) \times (0, 1)$$

such that

$$(4.1) \quad |x(\sigma(z))| < \frac{1}{2}\pi|z|^2 + \frac{\varepsilon}{2} \quad \text{for all } z \in D(a).$$


Figure 1 shows such an embedding. For details we refer to Lemma 3.1.8 of [42].

We claim that the symplectic embedding $\sigma \times \sigma : D(a) \times D(a) \rightarrow \mathbb{R}^4$ maps $B^4(a)$ to $\diamond(a + \varepsilon) \times (0, 1)^2$. Indeed, for $(z_1, z_2) \in B^4(a)$ we have $\pi(|z_1|^2 + |z_2|^2) < a$. Together with (4.1) we can estimate

$$\begin{aligned} |x_1((\sigma \times \sigma)(z_1, z_2))| + |x_2((\sigma \times \sigma)(z_1, z_2))| &= |x_1(\sigma(z_1))| + |x_2(\sigma(z_2))| \\ &< \frac{1}{2}(\pi|z_1|^2 + \pi|z_2|^2) + \varepsilon \\ &< \frac{a}{2} + \varepsilon, \end{aligned}$$

as claimed. $\square$

Later on, we shall often skip the $\varepsilon > 0$ appearing in the lemma, and simply think of the ball $B^4(a)$ as $\diamond(a) \times (0, 1)^2$.

FIGURE 1. The map σ .

4.2. Distorted diamonds. All of our embeddings, besides one, will start from a diamond $\diamond(a)$. For our full filling of $T(1, 1)$, however, we shall need to start from a *distorted diamond*.

Fix $a > 0$. Let $u_+ : [0, a] \rightarrow \mathbb{R}$ be a continuous and piecewise-linear function, with finitely many pieces, such that

$$u_+(0) = 0 \quad \text{and} \quad u'_+(\rho) \in [0, 1] \quad \text{on the linear pieces.}$$


Define the piecewise-linear function $u_- : [0, a] \rightarrow \mathbb{R}$ by $u_-(\rho) = u_+(\rho) - \rho$. Then $u_-(0) = 0$ and $u'_-(\rho) \in [-1, 0]$. Moreover, $u'_+(\rho) - u'_-(\rho) = 1$ on the linear pieces, and $u_+(a) - u_-(a) = a$. Let

$$\sigma_u : D(a) \rightarrow \left(u_-(a) - \frac{\varepsilon}{2}, u_+(a) + \frac{\varepsilon}{2}\right) \times (0, 1)$$

be a symplectic embedding such that

$$x(\sigma_u(z)) \in \left(u_-(\rho) - \frac{\varepsilon}{2}, u_+(\rho) + \frac{\varepsilon}{2}\right) \quad \text{for all } z \in D(a) \text{ with } \pi|z|^2 < \rho.$$

Figure 2 shows the image of concentric circles of the map σ_u for $a = 3$ and for the function $u_+ : [0, 3] \rightarrow \mathbb{R}$ with slope $\frac{1}{2}$ on $[0, \frac{3}{2}]$ and slope $\frac{1}{3}$ on $[\frac{3}{2}, 3]$.

FIGURE 2. The map σ_u for a function u_+ with two pieces.

Assume now that $a > 1$. Define d by $2d = a - 1$, and suppose that the functions u_+ , v_+ have only two pieces, with $u_+(\rho) = \frac{\rho}{2}$ for $\rho \in [0, 2d]$ and $v_+(\rho) = \frac{\rho}{2}$ for $\rho \in [0, 1]$. Computing as in the proof of Lemma 4.1 we find that the image of $B^4(a)$ under $\sigma_u \times \sigma_v$ is contained in $\diamond \times (0, 1)^2$, where $\diamond$ is as in Figure 3 (II). Again, we omitted $\varepsilon > 0$.




FIGURE 3. The diamond $\diamond(a)$, and a distorted diamond.

We call the factor $\diamond$ of such an image $\diamond \times (0, 1)^2$ a *distorted diamond*. A distorted diamond of size a therefore consists of a rectangle $(-d, d) \times (-\frac{1}{2}, \frac{1}{2})$ with $2d = a - 1$, a top and bottom triangle with base $2d$, the sum of whose heights is $v_+(a) - v_-(a) - 1 = a - 1 = 2d$, and two flaps with height 1, the sum of whose widths is $u_+(a) - u_-(a) - 2d = a - (a - 1) = 1$.

In fact one has the following more precise result.

Proposition 4.2. *Let $\diamond$ be a distorted diamond of size a . Then the product $\diamond \times (0, 1)^2$ is symplectomorphic to the open ball $B^4(a)$.*

Proof. In view of the above construction, for each compact subset $K \subset \diamond \times (0, 1)^2$ there exists $\hat{a} < a$ and a symplectic embedding $\hat{\varphi}: B^4(\hat{a}) \rightarrow \diamond \times (0, 1)^2$ such that $\text{Im } \hat{\varphi} \supset K$. The proposition thus follows from the following general result.

Lemma 4.3. *Let $V \subset \mathbb{R}^4$ be a bounded domain such that for each compact subset $K \subset V$ there exists $\hat{a} < a$ and a symplectic embedding $\hat{\varphi}: B^4(\hat{a}) \rightarrow V$ such that $\text{Im } \hat{\varphi} \supset K$. Then V is symplectomorphic to $B^4(a)$.*

Proof. Choose a sequence $K_1 \subset K_2 \subset \dots$ of compact subsets of V such that $\bigcup_j K_j = V$. Using the assumption of the lemma and the result from [30] that the space of symplectic embeddings of a closed ball into an open ball is connected, we construct a sequence $a'_1 < a_1 < a'_2 < a_2 < \dots$ with $a_j \rightarrow a$ and a sequence of symplectic embeddings $\varphi_j: B^4(a_j) \rightarrow V$ such that $\varphi_j(B^4(a'_j)) \supset K_j$ and $\varphi_{j+1}|_{B^4(a'_j)} = \varphi_j|_{B^4(a'_j)}$.

Define $\varphi: B^4(a) \rightarrow V$ by $\varphi(z) = \varphi_j(z)$ if $z \in B^4(a'_j)$. Then φ is a well-defined symplectic embedding. Moreover, $\varphi(B^4(a'_j)) = \varphi_j(B^4(a'_j)) \supset K_j$. Hence φ is onto $\bigcup_j K_j = V$. $\square$

In [44], Traynor proved Proposition 4.2 for the special case that $\diamond$ is the standard simplex $\{(x_1, x_2) \mid x_1, x_2 > 0, x_1 + x_2 < a\}$ by a different construction. Replacing in our construction the x_1 -coordinates by μx_1 and using the result from [35] that the space of symplectic embeddings of a closed four-dimensional ellipsoid into an open four-dimensional ellipsoid is connected, we find, more generally, that the product of a distorted diamond of size a , stretched along the x_1 -axis by μ , and the square $(0, 1)^2$, is symplectomorphic to the open ellipsoid $E(\mu a, a)$.

4.3. Shears. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function. Consider the x_1 -shear $\varphi(x_1, x_2) = (x_1 + f(x_2), x_2)$ of $\mathbb{R}^2$. Then the diffeomorphism

$$(4.2) \quad \widehat{\varphi}(x_1, x_2, y_1, y_2) = (x_1 + f(x_2), x_2, y_1, y_2 - f'(x_2)y_1)$$

is a symplectomorphism of $\mathbb{R}^4$. Indeed, this is just the “cotangent map”

$$(x_1, x_2, y_1, y_2) \mapsto \left(\varphi(x_1, x_2), (d\varphi(x_1, x_2))^T \right)^{-1}(y_1, y_2)$$

of the shear φ . We call a map $\widehat{\varphi}$ of the form (4.2) also an x_1 -shear. Similarly, an x_2 -shear $\varphi(x_1, x_2) = (x_1, x_2 + g(x_1))$ induces a symplectomorphism

$$(4.3) \quad (x_1, x_2, y_1, y_2) \mapsto (x_1, x_2 + g(x_1), y_1 - g'(x_1)y_2, y_2),$$

which we again call an x_2 -shear.

Let $U \subset \mathbb{R}^2(\mathbf{x}) := \mathbb{R}^2(x_1, x_2)$ be a domain, and consider the image $\widehat{\varphi}(U \times (0, 1)^2)$ of an x_1 -shear in $\mathbb{R}^2(\mathbf{x}) \times \mathbb{R}^2(\mathbf{y})$. This image fibers over $\varphi(U) \subset \mathbb{R}^2(\mathbf{x})$, with fiber $\{(y_1, y_2 - f'(x_2)y_1) \mid (y_1, y_2) \in (0, 1)^2\}$ over $\varphi(x_1, x_2)$, see Figure 4.




FIGURE 4. The fiber over $\varphi(x_1, x_2)$.

The projection $\pi_{\mathbf{y}}: \mathbb{R}^2(\mathbf{y}) \rightarrow \mathbb{R}^2(\mathbf{y})/\mathbb{Z}^2(1, 1)$ is injective on these fibers. It follows that if the projection $\pi_{\mathbf{x}}: \mathbb{R}^2(\mathbf{x}) \rightarrow \mathbb{R}^2(\mathbf{x})/\mathbb{Z}^2(\mu, 1)$ is injective, then also $\pi = \pi_{\mathbf{x}} \times \pi_{\mathbf{y}}: \widehat{\varphi}(U \times (0, 1)^2) \rightarrow T(\mu, 1)$ is injective. The same holds true for x_2 -shears. Moreover, if an x_1 -shear φ_1 and an x_2 -shear φ_2 are such that φ_2 fixes the set $\{\mathbf{x} \in \text{Im } \varphi_1 \mid \mathbf{x} \neq \varphi_1(\mathbf{x})\}$, and if $\pi_{\mathbf{x}}$ injects $\varphi_2 \circ \varphi_1(U)$ into $\mathbb{R}^2(\mathbf{x})/\mathbb{Z}^2(\mu, 1)$, then π injects $\widehat{\varphi}_2 \circ \widehat{\varphi}_1(U \times (0, 1)^2)$ into $T(\mu, 1)$.

We next give three examples illustrating the above embedding method.

Example 4.4. *A full filling of $T(2k^2, 1)$ for each $k \in \mathbb{N}$. We start from the diamond $\diamond(2k)$, with vertices $(\pm k, 0)$, $(0, \pm k)$. Using the linear shear $\varphi(x_1, x_2) = (x_1 + (2k - 1)x_2, x_2)$, the diamond is transformed into the rhomboid $R(k)$ with vertices $(\pm k, 0)$ and $\pm(2k^2 - k, k)$, see Figure 5.*



FIGURE 5. The rhomboid $R(k)$.

FIGURE 6. The image projects injectively.

The set $R(k)$ is a fundamental domain for the action of $\mathbb{Z}^2$ with generators $2k^2\partial_{x_1}$ and ∂_{x_2} . One way of seeing this is to translate the lower half of $R(k)$ by $2k^2\partial_{x_1} + k\partial_{x_2}$ to obtain the shape depicted in Figure 6. It follows from Proposition 4.2 that the ball $B^4(2k)$ symplectically embeds into $T(2k^2, 1)$.

Remark 4.5. Together with scaling and Remark 2.2, this gives an explicit full filling by one ball of $T(\mu, 1)$ for all $\mu = \frac{2m^2}{n^2}$ with m, n relatively prime. (Non-explicit full fillings of these tori follow from the computation of Seshadri constants (2.7), and from Proposition 1.6, together with Remark 2.2.)

Remark 4.6. By shears as in Example 4.4 one can also construct explicit full fillings for some special irrational tori, namely for those of the form $\mathbb{R}^4/\Lambda$ where the lattice splits as $\Lambda = \Lambda_x \times \Lambda_y$, such that Λ_y is the standard $\mathbb{Z}^2 \subset \mathbb{R}^2(\mathbf{y})$ and such that some linearly sheared diamond in $\mathbb{R}^2(\mathbf{x})$ is a fundamental domain for Λ_x .

Example 4.7. *Filling $\frac{8}{9}$ of $T(1,1)$.* Let $a = \frac{4}{3}$. The corners of the inner square of $\diamond(a)$ have coordinates $(\pm\frac{1}{3}, \pm\frac{1}{3})$. Choose f with $f(x_2) = 0$ for $-\frac{1}{3} \leq x_2 \leq \frac{1}{3}$ and $f'(x_2) = 1$ for $|x_2| > \frac{1}{3}$ (with rounding between). Then $f(-\frac{a}{2}) = -\frac{1}{3}$ and $f(\frac{a}{2}) = \frac{1}{3}$. The x_1 -shear induced by f then takes the upper triangle ABC in Figure 7 (I) to the shaded triangle $A'BC$, and similarly for the bottom triangle, while the rest of the diamond is untouched. The x_2 -shear induced by the function $-f$ then moves the left and right flaps of $\diamond(a)$ to the flaps shown in Figure 7 (II), while the rest of the image of the x_1 -shear is untouched. Therefore, a point in $\diamond(a)$ is affected by at most one of these two shears.

The composition of these two shears takes (a slight shrinking of) $\diamond(a)$ to the shaded domain in Figure 7 (II). This domain injects into $\mathbb{R}^2(\mathbf{x})/\mathbb{Z}^2$: It wraps up under the action of $\mathbb{Z}^2$ to a set covering all of the square $(-\frac{1}{2}, \frac{1}{2})^2$ besides the four black squares of area $(\frac{1}{6})^2$ each. For each $\varepsilon > 0$ we thus have constructed a symplectic embedding of a ball into $T(1,1)$ filling at least $\frac{8}{9} - \varepsilon$ of the volume of $T(1,1)$. This particular method does not do better because we need A and B to be vertically equidistant from the line $x_2 = \frac{1}{2}$ to do the x_1 -shear so symmetrically.





FIGURE 7. Filling $\frac{8}{9}$ of $T(1,1)$.


Example 4.8. *Filling $\frac{49}{50}$ of $T(1,1)$.* Let $a = \frac{7}{5}$. The idea is to divide the square representing $T(1,1)$ into two rectangles, one the maximum rectangle of height 1 that lies in the diamond $\diamond(a)$ (and hence has width $a - 1 = \frac{2}{5}$), and the other of width $2 - a = \frac{3}{5}$, see Figure 8 (I).


We shear the top triangle by a strong x_1 -shear to the left, the bottom triangle by a strong x_1 -shear to the right, and then shear the flaps by x_2 -shears in a symmetric way so as to free

FIGURE 8. Filling $\frac{49}{50}$ of $T(1,1)$, schematically.

triangles into which the sheared top and bottom triangles fit, when projected to the torus. The freed triangles have height $\frac{1}{5}$ and width $\frac{1}{2}$, while the triangles fitting in have the same height, but width $\frac{2}{5}$ only. In Figure 8 (II), one sees the image of the top triangle and its translate by $\partial_{x_1} - \partial_{x_2}$, the image of the bottom triangle and its translate by ∂_{x_2} , as well as the image of the left flap and its translate by ∂_{x_1} .

To make this construction precise, fix a small $\varepsilon > 0$, and decompose the diamond $\diamond(a - \varepsilon)$ into four triangles and a rectangle of height 1 and width $\frac{2}{5}$ from each of whose four vertices a simplex of width $\frac{\varepsilon}{2}$ has been removed. For notational convenience, we also translate $\diamond(a - \varepsilon)$ by $\frac{1}{2}\partial_{x_2}$ (see Figure 9).

FIGURE 9. The decomposition of the diamond $\diamond(a - \varepsilon)$.

FIGURE 10. Filling $\frac{49}{50}$ of $T(1,1)$.

The x_1 -shear, that moves the top triangle to the left and the bottom triangles to the right, has support in $\{x_2 > 1 - \frac{\varepsilon}{4}\} \cup \{x_2 < 0\}$. Near the upper triangle, this shear is very strong on $\{1 - \frac{\varepsilon}{4} < x_2 < 1\}$, and up to a minor shear, it is a translation on $\{1 < x_2 < \frac{a-\varepsilon}{2} + \frac{1}{2}\}$. It fixes the points $A = (\frac{\varepsilon}{4}, 1 - \frac{\varepsilon}{4})$ and $B = (\frac{2}{5} - \frac{\varepsilon}{4}, 1 - \frac{\varepsilon}{4})$, but translates $C = (\frac{\varepsilon}{2}, 1)$ to $C' = (-\frac{2}{5} + \frac{\varepsilon}{2}, 1)$ and $D = (\frac{2}{5} - \frac{\varepsilon}{2}, 1)$ to $D' = (-\frac{\varepsilon}{2}, 1)$. On the bottom triangle, the x_1 -shear acts very strongly on $\{-\frac{\varepsilon}{4} < x_2 < 0\}$. It fixes the points $X = (\frac{\varepsilon}{2}, 0)$ and $Y = (\frac{2}{5} - \frac{\varepsilon}{2}, 0)$, but translates $W = (\frac{3}{4}\varepsilon, -\frac{\varepsilon}{4})$ to $W' = (\frac{2}{5} + \frac{\varepsilon}{2}, -\frac{\varepsilon}{4})$ and $Z = (\frac{2}{5} - \frac{3}{4}\varepsilon, -\frac{\varepsilon}{4})$ to $Z' = (\frac{4}{5} - \varepsilon, -\frac{\varepsilon}{4})$.

In Figure 10 we drew the projection of the (dark) top triangle $\{1 - \frac{\varepsilon}{4} < x_2\}$ to the fundamental domain $(0,1) \times (0,1)$ of the usual $\mathbb{Z}^2$ -action, but we did not draw the projection of the bottom triangle $\{x_2 < 0\}$. In order to see that the image projects injectively to $T(1,1)$, notice that the point C lies strictly above the segment $\overline{D'B}$, that W' lies on the right of B , and that B lies $\frac{\varepsilon}{4}$ below C and W' lies $\frac{\varepsilon}{4}$ below X . Therefore, the translate by ∂_{x_2} of the segment $\overline{XW'}$ lies above the segment $\overline{D'B}$.

5. PROOF OF THEOREM 1

Since we already proved Theorem 1.9, it only remains to treat the product torus $T(1, 1)$. So let $a = \sqrt{2}$. We want to find, for each $\varepsilon > 0$, a symplectic embedding of the ball $B^4(a - \varepsilon)$ into the torus $T(1, 1)$. We describe the schematic embedding for $\varepsilon = 0$. From this, an actual embedding for $\varepsilon > 0$ is obtained exactly as in Example 4.8.

As in that example, given the diamond $\diamond(a)$, we are led to decompose the square $(0, 1) \times (-\frac{1}{2}, \frac{1}{2})$ into two rectangles, and to try to fill the right rectangle $(a - 1, 1) \times (-\frac{1}{2}, \frac{1}{2})$ with the four triangles, see Figure 11 (I).




FIGURE 11. The decomposition of $\diamond(a)$, and its distortion.

The width of this rectangle is $1 - (a - 1) = 2 - \sqrt{2} =: \frac{1+b}{2}$, with $b := 3 - 2\sqrt{2}$. The new idea is to start from a distorted diamond, instead, whose left flap has width $\frac{1+b}{2}$, and so will just fit into the right rectangle. The right flap of the distorted diamond then has width $\frac{1-b}{2}$. The height h_t of its top triangle will be chosen later. The bottom triangle then has height $h_b = 2d - h_t$, where $2d = a - 1 = \sqrt{2} - 1$. Note that $2d = \frac{1-b}{2}$. After a translation, the distorted diamond then looks as in Figure 11 (II).

We put the two flaps into the rectangle $(0, \frac{1+b}{2}) \times (0, 1)$ so as to create *triangular* top and bottom regions R_- and R_+ , both with a base of width $\frac{1-b}{2}$, and their tips at distance b from the closest edge, see Figure 12. There is a unique way to do this if we require the tip of the right flap to lie on the lower left corner of the rectangle. We realize this position of the flaps of the distorted diamond by x_2 -shears (where in the figure the image of the left flap is translated by ∂_{x_1}), by first putting in the dark flap flush with the bottom, then putting in the light flap flush with the image of the dark flap, and finally shearing the right part of the flaps up in parallel to make the top triangle narrower (of width $\frac{1-b}{2}$) and simultaneously creating the bottom triangle of the same width. Define h_t to be the height of the lower triangle R_- . Then the height h_b of the upper triangle R_+ necessarily satisfies $\frac{1-b}{2} = h_t + h_b$. Indeed, the area of the distorted diamond is 1, and so the sum of the area of its top and bottom triangles




FIGURE 12. Filling the right rectangle with the distorted triangles.

must equal the sum of the area of R_- and R_+ ; since all four triangles have base $\frac{1-b}{2}$, the sum of their heights must be equal too.

To check, the slope s_1 in Figure 12 is $\frac{1}{(1+b)/2} = \frac{2}{1+b}$. Therefore, $h_t = bs_1 = \frac{2b}{1+b}$. Similarly, $s_2 = s_1 + \frac{2}{1-b} = \frac{4b}{(1-b)(1+b)}$, and hence $h_b = \frac{4b^2}{(1-b)(1+b)}$. Since $b = 3 - 2\sqrt{2}$, one indeed finds $h_t + h_b = \frac{1-b}{2}$.

As in Example 4.8, we can therefore use an x_1 -shear to bring the top triangle of the distorted diamond into R_- and the bottom triangle into R_+ . $\square$

6. PROOFS OF COROLLARIES 1.2 AND 1.3

Proof of Corollary 1.2. Consider the 4-torus T^4 and its blow-up $X_1 = T^4 \# \overline{\mathbb{C}P^2}$. Fix an orientation of X_1 . Denote by $E \in H_2(X_1; \mathbb{Z})$ the homology class of the exceptional divisor (with some orientation) in X_1 . We need to show that the symplectic cone of X_1 is

$$(6.1) \quad \mathcal{C}(X_1) = \{ \alpha \in H^2(X_1; \mathbb{R}) \mid \alpha^2 > 0, \alpha(E) \neq 0 \}.$$

We first prove the inclusion $\subset$ in (6.1). The condition $\alpha^2 > 0$ holds because α is represented by a symplectic form compatible with the given orientation of X_1 . The condition $\alpha(E) \neq 0$ follows from Taubes' work on the relation between Seiberg–Witten and Gromov invariants, according to which for any symplectic form ω on X_1 the class E is representable by an embedded sphere on which ω is non-degenerate, see [45, 46].

We now prove the inclusion $\supset$ in (6.1). The projection $\pi: X_1 \rightarrow T^4$ induces an orientation on T^4 . The classes in $H^2(X_1; \mathbb{R})$ can be written as $\pi^*\beta - ae$ where $\beta \in H^2(T^4; \mathbb{R})$ and $a \in \mathbb{R}$, and where $e = \text{PD}(E)$ is the Poincaré dual of E . Since $e^2 = -1$, the set on the right hand side of (6.1) becomes

$$\{\pi^*\beta - ae \in H^2(X_1; \mathbb{R}) \mid \beta \in H^2(T^4; \mathbb{R}), \beta^2 > a^2 > 0\}.$$

Fix $\beta \in H^2(T^4; \mathbb{R})$ and $a > 0$ with $\beta^2 > a^2 > 0$. Since $\beta^2 > 0$, we can represent β by a linear symplectic form on T^4 compatible with the given orientation. Since $\beta^2 > a^2 > 0$, Theorem 1.1 guarantees a symplectic embedding of $B^4(a)$ into (T^4, ω) . The symplectic form on the corresponding symplectic blow-up of (T^4, ω) represents the class $\pi^*\beta - a'e$, where either $a' = a$ or $a' = -a$.

There exists an orientation-preserving diffeomorphism $\varphi: X_1 \rightarrow X_1$ which acts on $H_2(X_1; \mathbb{Z})$ by sending E to $-E$ while fixing the orthogonal complement $\pi^*(H^2(T^4; \mathbb{R}))$ of E (with respect to the intersection pairing). In particular, $\varphi^*(\pi^*\beta - a'e) = \pi^*\beta + a'e$. Therefore, both $\pi^*\beta - ae$ and $\pi^*\beta + ae$ belong to the symplectic cone $\mathcal{C}(X_1)$. $\square$

Proof of Corollary 1.3. Throughout this proof, we denote by ω_d a linear symplectic form on T^4 such that (T^4, ω_d) is symplectomorphic to $T(1, d)$. By Corollary 1.2, the class $\alpha(a) = \pi^*([\omega_1]) - a \text{PD}(E)$ admits a symplectic representative $\omega_{1,a}$ for all $0 < a < \sqrt{2}$. If there were a Kähler representative of $\alpha(a)$, then we could perform the blow down in a Kähler way to obtain a Kähler structure on T^4 of symplectic type $(1, 1)$ which contains a symplectically and holomorphically embedded ball $B^4(a)$. Since any such Kähler structure corresponds to a principally polarized abelian surface, we would conclude, together with Remark 2.5, that its Seshadri constant is at least a . This contradicts Steffens' estimate (2.4) whenever $a > \frac{4}{3}$.

More generally, we can start from the torus $T(d, 1)$ with $d \neq 2k^2$ an integer. By (2.8), there exists a non-empty interval $(\frac{\varepsilon_d}{\sqrt{d}}, \sqrt{2d})$ such that for a in this interval, the class $\pi^*([\omega_d]) - a \text{PD}(E)$ on X_1 admits a symplectic representative but no Kähler representative.

Let now $k \geq 2$, and choose $a \in (\frac{4}{3}, \sqrt{2})$. For $\varepsilon_2, \dots, \varepsilon_k > 0$ sufficiently small, we can symplectically blow up $(X_1, \omega_{1,a})$ by weights $\varepsilon_2, \dots, \varepsilon_k$, and obtain a symplectic representative of the class $\pi_k^*([\omega_1]) - a \text{PD}(E) - \sum_{j=2}^k \varepsilon_j \text{PD}(E_j)$. This class does not contain a Kähler representative. Otherwise, we could blow-down along the exceptional divisors $E_2, \dots, E_k$ to obtain a Kähler representative of the class $\pi^*([\omega_1]) - a \text{PD}(E)$ on X_1 . $\square$

7. REMARKS AND QUESTIONS

1. Symplectic forms on T^4 . We have worked throughout with a linear symplectic form on T^4 . It is not known whether every symplectic form on T^4 is isotopic to a linear form, or even whether it is symplectomorphic to such a form.

2. Filling $T(1, 1)$. The general filling methods of §3 work for all linear tori except the “square torus” $T(1, 1)$. The most interesting remaining question here is: can $T(1, 1)$ be fully filled by $E(1, b)$ for every b ? By Theorem 1.10 and because ellipsoids are nested, it would suffice to prove this for a dense set of $b \in (1, 2)$; cf. [40, Lemma 1.1]. If we could extend the explicit construction of §5 to a full filling of the tori $T(b, 1)$, $b \in (1, 2)$, by a ball, then, by rescaling the x_1 -coordinates of the domain and target by a factor of $\frac{1}{b}$ one would obtain a full filling of $T(1, 1)$ by a dilate of $E(\frac{1}{b}, 1) = \frac{1}{b}E(1, b)$. However the construction in §5 is very delicate and there seems no easy way to extend it to other $T(b, 1)$.

3. Very full fillings. There is a stronger version of full filling: rather than asking whether one can fill an arbitrarily large fraction of the volume of a manifold M with a ball, one could ask whether M has a set of full measure that is symplectomorphic to an open ball. In other words, if $a = c_G(M, \omega)$ does the open ball $B^4(a)$ embed symplectically in M ? Let us say that in this case (M, ω) has a very full filling (by one ball). (There are similar versions for other filling problems.) When a rational or ruled manifold has a full filling, it also has a very full filling because one can argue as in the proof of Lemma 4.3, using the fact that in this case the space of ball embeddings is connected. However, these general arguments do not apply to tori, and it is unclear whether $T(1, 1)$, for example, has a very full filling by one ball. On the other hand, the explicit fillings in Example 4.4 and Remark 4.5 give very full fillings.

4. The isotopy problem. For some symplectic four-manifolds (M, ω) , such as the complex projective plane or a product of 2-spheres, it is known that the space of symplectic embeddings of a given (closed) ball into (M, ω) is connected, see [30, 33]. For tori, this is a completely open problem. For many balls $B^4(a)$, our embedding constructions yield various symplectic embeddings into tori $T(\mu, 1)$, for which we do not know whether they are symplectically isotopic.

As a first example, consider, for some fixed small $\varepsilon > 0$, the symplectic embeddings of a ball filling $\frac{8}{9} - \varepsilon$ of $T(1, 1)$ that are illustrated, for $\varepsilon = 0$, in Figure 13. Here, the embedding $(+-)$ is the one of Example 3 in Section 4.3, and the other three embeddings are obtained in the same way. Are these balls symplectically isotopic in $T(1, 1)$? Note that, for instance, the (not Hamiltonian) symplectomorphism $(x_1, y_1, x_2, y_2) \mapsto (-x_1, -y_1, x_2, y_2)$ of $T(1, 1)$ maps the ball $(++)$ to $(--)$ and maps $(+-)$ to $(-+)$.





FIGURE 13. Four embeddings of $B^4(\frac{4}{3})$ into $T(1, 1)$.

As a second example, consider the following two full fillings of $T(\frac{9}{8}, 1)$: The first filling is the one obtained from the explicit full filling of $T(1, 72)$ via Lemma 2.1. The second filling is similar to the embedding in Example 3 of Section 4.3. We decompose the diamond $\diamond(\frac{3}{2})$ and the rectangle $(0, \frac{9}{8}) \times (0, 1)$ as in Figure 14 (I), and shear the triangles as shown in Figure 14 (II).⁶ These two embeddings are clearly different. Are they symplectically isotopic?

More generally, it is not known whether there is any $\varepsilon \in (0, \sqrt{2})$ such that the space of symplectic embeddings of $B^4(\varepsilon)$ into $T(1, 1)$ is connected.

⁶In fact, this is the first in a family of full fillings of $T(\frac{(2k+1)^2}{2(k+1)^2}, 1)$, $k \geq 1$, by the diamond $\diamond(\frac{2k+1}{k+1})$ in which the top triangle $x_2 \geq \frac{1}{2}$ is sliced into k horizontal slices of heights $\frac{1}{(k+1)^2}, \frac{2}{(k+1)^2}, \dots, \frac{k}{(k+1)^2}$ and then sheared to the right so that the right edge of the j th piece lies over an x_1 -interval of length $\frac{2j-1}{2(k+1)^2}$, while the bottom triangle is sheared symmetrically to the left.

FIGURE 14. Another full filling of $T(\frac{9}{8}, 1)$.

5. Uniqueness of symplectic structures on the blow-up of T^4 . Suppose that ω_1, ω_2 are two cohomologous symplectic forms on the blow-up of a given symplectic 4-manifold (M, ω) that are obtained by blow-up from two ball embeddings into M . Then it is shown in [39, Prop. 7.20] that ω_1, ω_2 are isotopic if and only if the two ball embeddings are symplectically isotopic. Moreover, the easiest way to show that two blow-ups of some symplectic 4-manifold are symplectomorphic is to use this equivalence. (Of course sometimes, as with the $(++)$, $(--)$ embeddings mentioned in 4. above, there are obvious symplectomorphisms that take one embedding to another and hence one blow-up to the other.) Since the symplectic isotopy problem for embeddings of balls in 4-dimensional tori is open, the uniqueness problem for symplectic forms on their blow-ups is open too, even if we restrict consideration to forms on the blow-up that blow down to linear forms.

6. Higher dimensions. The filling methods used in §3 work only in dimension 4. Although many of the explicit arguments in §4 extend to higher dimensions, the higher dimensional analogs of the diamond $\diamond$ (e.g. the octahedron) do not tile Euclidean space. Therefore there seem to be no simple explicit full fillings of tori by balls or ellipsoids in higher dimensions along the lines of Example 4.4. As we explained in Section 2.3, one can get some (presumably rather weak) lower bounds for the ball filling number of tori of dimension $2n \geq 6$ from the computations of Seshadri constants in [3] and [12]. For example, when $n = 3, 4$ we have:

$$p(T^6) \geq \frac{288}{343} \quad \text{and} \quad p(T^8) \geq \frac{2}{3}.$$

It is not clear how to do better than this, or even how to realize these bounds by explicit embeddings. It is also not clear how to find explicit embeddings in dimension 4 that do better than some of the sharper Seshadri constants, for example $\frac{360}{361}$ for $T(1, 5)$.

REFERENCES

- [1] T. Bauer. Seshadri constants and periods of polarized abelian varieties. With an appendix by the author and T. Szemberg. *Math. Ann.* **312** (1998) 607–623.
- [2] T. Bauer. Seshadri constants on algebraic surfaces. *Math. Ann.* **313** (1999) 547–583.
- [3] T. Bauer and T. Szemberg. Local positivity of principally polarized abelian threefolds. *J. Reine Angew. Math.* **531** (2001) 191–200.

- [4] P. Biran. Symplectic packing in dimension 4. *Geom. Funct. Anal.* **7** (1997) 420–437.
- [5] P. Biran. A stability property of symplectic packing. *Invent. Math.* **136** (1999) 123–155.
- [6] P. Biran and K. Cieliebak. Symplectic topology on subcritical manifolds. *Comment. Math. Helv.* **76** (2001) 712–753.
- [7] C. Birkenhake and H. Lange. *Complex abelian varieties*. Grundlehren der mathematischen Wissenschaften, vol. 302, 2nd edition, Springer-Verlag, Berlin, 2004.
- [8] N. Buchdahl. On compact Kähler surfaces. *Ann. Inst. Fourier* **49** (1999) 287–302.
- [9] O. Buse and R. Hind. Symplectic embedding of ellipsoids in dimension greater than four. Preprint 2011.
- [10] O. Buse and M. Pinsonnault. Packing numbers of rational ruled 4-manifolds. arXiv:1104.3362
- [11] F. Catanese, K. Oguiso and T. Peternell. On volume preserving complex structures on real tori. *Kyoto J. Math.* **50** (2010) 753–775.
- [12] O. Debarre. *Higher-dimensional algebraic geometry*. Universitext, Springer-Verlag, New York, 2001.
- [13] T. Drăghici. The Kähler cone versus the symplectic cone. *Bull. Math. Soc. Sci. Math. Roumanie* **42** (1999) 41–49.
- [14] G. Elenicwajg and O. Forster. Vector bundles on manifolds without divisors and a theorem on deformations. *Ann. Inst. Fourier* **32** (1982) 25–51.
- [15] M. Gromov. Pseudo-holomorphic curves in symplectic manifolds. *Invent. math.* **82** (1985) 307–347.
- [16] H. Hofer and E. Zehnder. *Symplectic Invariants and Hamiltonian Dynamics*. Birkhäuser, Basel, 1994.
- [17] M. Hutchings. Quantitative embedded contact homology [new] arXiv:1005.2260. To appear in *Journal of Differential Geometry*.
- [18] M.-Y. Jiang. Symplectic embeddings from $\mathbb{R}^{2n}$ into some manifolds. *Proc. Roy. Soc. Edinburgh Sect. A* **130** (2000) 53–61.
- [19] Y. Karshon and S. Tolman. The Gromov width of complex Grassmannians. *Algebr. Geom. Topol.* **5** (2005) 911–922.
- [20] J. L. Lagrange. Solution d’un Problème d’Arithmétique. *Oeuvres*, tome 1, 671–732.
- [21] F. Lalonde and D. McDuff. The geometry of symplectic energy. *Ann. of Math.* **141** (1995) 349–371.
- [22] F. Lalonde and D. McDuff. Hofer’s L^∞ -geometry: energy and stability of Hamiltonian flows, part II. *Invent. math.* **122** (1995) 35–69.
- [23] F. Lalonde and D. McDuff. The classification of ruled symplectic 4-manifolds. *Math. Res. Lett.* **3** (1996) 769–778.
- [24] A. Lamari. Le cone Kählérien d’une surface. *J. Math. Pures Appl.* **78** (1999) 249–263.
- [25] R. Lazarsfeld. Lengths of periods and Seshadri constants of abelian varieties. *Math. Res. Lett.* (1996) 439–447.
- [26] R. Lazarsfeld. *Positivity in algebraic geometry. I*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, vol. 48, Springer-Verlag, Berlin, 2004.
- [27] T. J. Li and A. Liu. Symplectic structure on ruled surfaces and a generalized adjunction formula. *Math. Res. Lett.* **2** (1995) 453–471.
- [28] T.-J. Li and M. Usher. Symplectic forms and surfaces of negative square. *J. Symplectic Geom.* **4** (2006) 71–91.
- [29] G. Lu. Gromov-Witten invariants and pseudo symplectic capacities. *Israel J. Math.* **156** (2006), 1–63.
- [30] D. McDuff. Blowups and symplectic embeddings in dimension 4. *Topology* **30** (1991) 409–421.
- [31] D. McDuff. The structure of rational and ruled symplectic 4-manifolds. *J. Amer. Math. Soc.* **3** (1990) 679–712.
- [32] D. McDuff. Notes on ruled symplectic 4-manifolds. *Trans. Amer. Math. Soc.* **345** (1994) 623–639.
- [33] D. McDuff. From symplectic deformation to isotopy. Topics in symplectic 4-manifolds (Irvine, CA, 1996), 85–99, *First Int. Press Lect. Ser., I*, Int. Press, Cambridge, MA, 1998.
- [34] D. McDuff. Geometric variants of the Hofer norm. *J. Symplectic Geom.* **1** (2002) 197–252.
- [35] D. McDuff. Symplectic embeddings of 4-dimensional ellipsoids. *J. Topol.* **2** (2009) 1–22.
- [36] D. McDuff. Symplectic embeddings and continued fractions: a survey. arXiv:0908.4387, *Journ. Math. Soc. Jap.* **4** (2009) 121–139.
- [37] D. McDuff. The Hofer conjecture on embedding symplectic ellipsoids. arXiv:1008.1885. *Journal of Differential Geometry* **88** (2011) 519–532.
- [38] D. McDuff and L. Polterovich. Symplectic packings and algebraic geometry. *Invent. math.* **115** (1994) 405–429.

- [39] D. McDuff and D. Salamon. *Introduction to symplectic topology*. Second edition. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1998.
- [40] D. McDuff and F. Schlenk. The embedding capacity of 4-dimensional symplectic ellipsoids. arXiv:0912.0532. To appear in *Annals of Mathematics*.
- [41] D. McDuff and J. Slimowitz. Hofer-Zehnder capacity and length minimizing Hamiltonian paths. *Geom. Topol.* **5** (2001) 799–830.
- [42] F. Schlenk. *Embedding problems in symplectic geometry*. de Gruyter Expositions in Mathematics **40**. Walter de Gruyter Verlag, Berlin, 2005.
- [43] A. Steffens. Remarks on Seshadri constants. *Math. Z.* **227** (1998) 505–510.
- [44] L. Traynor. Symplectic packing constructions. *J. Differential Geom.* **41** (1995) 735–751.
- [45] C. H. Taubes. The Seiberg–Witten and Gromov invariants. *Math. Res. Lett.* **2** (1995) 221–238.
- [46] C. H. Taubes. *Seiberg–Witten and Gromov Invariants in Symplectic 4-manifolds*. First International Press Lecture Series, 2. International Press, Somerville, MA, 2000.

(J. LATSCHEV) FACHBEREICH MATHEMATIK, UNIVERSITÄT HAMBURG, BUNDESSTRASSE 55, 20146 HAMBURG, GERMANY

E-mail address: `janko.latschev@math.uni-hamburg.de`

(D. MCDUFF) DEPARTMENT OF MATHEMATICS, BARNARD COLLEGE, COLUMBIA UNIVERSITY, NEW YORK, NY 10027-6598, USA.

E-mail address: `dmcduff@barnard.edu`

(F. SCHLENK) INSTITUT DE MATHÉMATIQUES, UNIVERSITÉ DE NEUCHÂTEL, RUE ÉMILE ARGAND 11, CP 158, 2000 NEUCHÂTEL, SWITZERLAND

E-mail address: `schlenk@unine.ch`